### Coherent structures in Quantum Turbulence

(Waves are present...but not discussed)

Andrew Baggaley (Newcastle) Jason Laurie (Aston)

### Classical (viscous) turbulence

- In a 3D classical turbulent flow, large scale eddies break up into smaller eddies, these into smaller ones and so on...(Richardson Cascade)
- If there is a large inertial range between the forcing and dissipation scale (i.e. high Re) then the flow of energy through scales is characterized by a constant energy flux .
- Dimensional analysis leads to a power-law scaling for the energy spectrum,

$$E(k) = C\epsilon^{2/3}k^{-5/3}$$

$$\frac{\partial \mathbf{v}}{\partial t} + (\mathbf{v} \cdot \nabla)\mathbf{v} = -\frac{1}{\rho}\nabla p + \nu\nabla^2 \mathbf{v}$$

$$v = \sin(x) \Rightarrow v \frac{\partial v}{\partial x} \sim \sin(2x)$$



## Classical Vorticity







### Quantum Fluids

- Atom with momentum p=mv has wavelength  $\lambda=h/p$
- Average kinetic energy  $mv^2/2 \approx k_B T$
- Wavelength increases with decreasing T:

$$\lambda \approx rac{h}{\sqrt{mk_B T}}$$

• Compare  $\lambda$  against the average distance between atoms, d:



BEC occurs when  $\lambda \approx d$ 

$$\Gamma = \oint_C \mathbf{v} \cdot d\mathbf{l} \in \mathbb{R}$$



$$\Gamma = \oint_C \mathbf{v} \cdot d\mathbf{l} = \frac{2\pi\hbar}{m}n$$



Paoletti et al., 2008

### Kuchemann:

#### "

vortices are the sinews and muscles of fluid motions"





These two drawings from Leonardo's studies in hydrodynamics, now in the collection of the Library of the Institut de France, represent vortex formation with the flow of water around an obstacle or through an opening in a partition within a trough. The second figure of symmetric counter-rotating vortices brings to mind Theodore von Karman's vortex street of asymmetric counter-rotating vortices formed in the wake of a circular cylinder moving through a field. In his 1954 Aerodynamics, von Karman wrote: "I do not claim to have discovered these vortices: they were known long before I was born. The earliest picture in which I have sen them is one in a church in Bologna, Italy, where St. Christopher is shown carrying the child Jesus across a flowing stream. Behind the saint's naked foot the painter indicated alternating writices."

## If this is true then Quantum Turbulence represents the 'skeleton'



### Yet we still see 'classical' behaviour



## Coherent structures

- In classical turbulence vorticity is concentrated in vortical 'worms' (She & al, Nature, 1990; Goto, JFM, 2008)
- Are there vortex bundles in quantum turbulence ?
- Would allow a mechanism for vortex stretching, i.e. stretch the bundle.



 $\frac{D\boldsymbol{\omega}}{Dt} = (\boldsymbol{\omega}\cdot\nabla)\mathbf{v} + \nu\nabla^2\boldsymbol{\omega}$ 

## Mathematical approach

### 3 distinct scales/numerical approaches







Barenghi et al. (2014)

### Gross-Pitaevskii



### Point Vortex/VFM



Course-Grained HVBK





### Mutual friction



## $\frac{d\mathbf{s}}{dt} = \mathbf{v}_s^{\text{tot}} + \alpha \mathbf{s}' \times \left(\mathbf{v}_n^{\text{ext}} - \mathbf{v}_s^{\text{tot}}\right) - \alpha' \mathbf{s}' \times \left[\mathbf{s}' \times \left(\mathbf{v}_n^{\text{ext}} - \mathbf{v}_s^{\text{tot}}\right)\right]$

Normal viscous fluid coupled to inviscid superfluid via mutual friction.

Superfluid component extracts energy from normal fluid component via Donelly-Glaberson instability, amplification of Kelvin waves.

Kelvin wave grows with amplitude:

 $\mathcal{A}(t) = \mathcal{A}(0)e^{\sigma t}$ 

$$\sigma(k) = \alpha(kV - \nu'k^2)$$

Counterflow Turbulence



# Generation of bundles at finite temperatures



VGatesslanckingna Milonioisv Kidpkik-& Ronuseda, PRB, 20008

### Reconnections: Bundles remain intact





Numerical simulations using both GPE and vortex filament method.









Alamri *et al.* 2008

## Decomposition of a tangle





AWB, Laurie & Barenghi, 2012

### Motivation







#### Roussel, Schneider & Farge, 2005

## Numerical results





Left, frequency spectra (red polarised ; black total), right energy spectrum, upper random component, lower polarised component.

## Experimental detection

Rusaouen et al., 2017



Presence of coherent structures inferred from intermittent pressure drops



### Hall-Vinen-Bekarevich-Khalatnikov Equations

Course-grained, macroscopic model

$$\begin{split} &\frac{\partial \mathbf{v}_n}{\partial t} + \left(\mathbf{v}_n \cdot \nabla\right) \mathbf{v}_n = -\frac{1}{\rho} \nabla P + \mu \nabla^2 \mathbf{v}_n + \frac{\rho_s}{\rho} \mathbf{F}, \quad \nabla \cdot \mathbf{v}_n = 0, \\ &\frac{\partial \mathbf{v}_s}{\partial t} + \left(\mathbf{v}_s \cdot \nabla\right) \mathbf{v}_s = -\frac{1}{\rho} \nabla P - \frac{\rho_n}{\rho} \mathbf{F}, \quad \nabla \cdot \mathbf{v}_s = 0. \\ &\mathbf{F} \simeq \alpha \rho_s \langle |\omega_{\mathbf{s}}| \rangle (\mathbf{v}_s - \mathbf{v}_n) \end{split}$$

$$\rho_s \gg \rho_n: \qquad \nabla^2 P \sim \frac{\rho_s}{2} (\omega_s^2 - \sigma_s^2)$$



## A single bundle in isolation





## Turbulent Tangle

$$\hat{F}(|\mathbf{k}|) = \exp\left(-\frac{|\mathbf{k}|^2}{2k_f^2}\right) \qquad k_f = 2\pi/l_f$$







### Summary

- Coherent vortical structures are present in the quasiclassical regime of Quantum Turbulence.
- Important (essential?) for K41 like statistical properties of QT.
- Good agreement between macroscopic HVBK model and mesoscale vortex approach.
- Interesting high pressure signal found in the Vinen regime.

The End